

Well + *Easy*

A WELL + EASY BOOK

The Whole GI Way

Keep the foods you love.
Know what to do today.
Adapt simply when life changes.

A thoughtful guide to caring for your blood sugar, weight, and strength - and living the life you want to keep living.

By Jen
Founder of Well + Easy

Make yourself comfortable

Read a little, stay awhile, come back whenever you like. The contents and research notes in this edition are clickable. Web links lead to further reading and Well + Easy resources.

Revised edition · 2026

© 2026 Well + Easy

Whole GI brings clinically studied principles into a framework developed and refined with clients over thirteen years. This book helps you understand the practices, use them in daily life, and notice the benefits. Educational guidance complements your individual medical care.

Read the web edition at wellandeasy.com/whole-gi-way.

Contents

A place to begin	5
1. The care you have already been giving	7
2. What do you want your health for?	9
3. Blood sugar, without the blame	11
4. How a food ranking became a kitchen question	13
5. What the evidence can offer	15
6. Keep the foods you love	17
7. Look at the whole meal	19
8. Preparation, portions, and a little perspective	22
9. A rhythm for an ordinary day	24
10. Breakfast you can recognize	28
11. Give lunch a proper place	30
12. Dinner can be simple and still be dinner	32
13. Snacks, dessert, and the end of earning food	35
14. When today has less to give	38
15. A small appetite still deserves nourishment	40
16. Strength belongs in the conversation	42

17. A changing body deserves a changing plan	44
18. Beyond the plate, without another project	46
19. One table, different needs	49
20. Restaurants, holidays, and a full life	51
21. A kitchen that helps you on Tuesday	53
22. When a difficult week becomes part of the way	55
23. The first thirty days, gently	57
24. Support you can keep close	59
25. The life on the other side of the kitchen	61
Notes and further reading	63
Keep exploring	68

WELCOME

A place to begin

Perhaps you have a cup of something warm beside you. Perhaps you are reading this on your phone while dinner is doing whatever dinner is doing. Perhaps you came looking for a straightforward answer about blood sugar and found yourself opening a book.

Welcome. There is room for you here.

You do not need to arrive with a clean refrigerator, a recent grocery trip, or an impressive history of sticking with things. You do not need to promise that this time will be different. You can bring the uncertainty, the favorite foods, the medical appointment you are still thinking about, and the very reasonable wish that taking care of yourself could feel a little less complicated.

This book begins with three possibilities: you can keep foods you love; you can learn what to do today; and you can adapt when life changes. Together, those possibilities offer a way to care for your health without making food the center of every waking hour.

Whole GI is my practical framework for bringing glycemic index, glycemic load, meal composition, preparation, timing, and the circumstances of an actual day into the same conversation. The work behind it began in 2013. For thirteen years, I have tested and refined this framework with clients: teaching it, listening to what changes, and finding ways to make the science useful in an ordinary kitchen. Whole GI brings clinically studied nutrition principles together in a client-tested way of living.

Much of that listening has been to women in their fifties, sixties, seventies, and beyond. Women who already put care into their food. Women who have been told to watch their blood sugar without being given much help with breakfast tomorrow. Their experiences are woven into the questions this book asks.

Men belong here, too. So do younger adults, people feeding a partner, people living alone, and people helping someone they love. There is no single household, body size, appetite, or age that earns a place at this table. What matters is that you would like a practical way to feed yourself with care.

You can read from beginning to end or begin with the chapter that meets your day. The science chapters explain the foundations. The kitchen chapters help you use them. The later chapters make room for the weeks when your energy, appetite, or plans change. Small pauses invite you to reflect; you never have to write an answer.

Examples in these pages are teaching situations, rather than accounts of particular clients. Meal ideas are flexible illustrations, not individualized prescriptions. Research notes are numbered so you can follow a thought into the evidence when you want to. You are equally welcome to stay with the story.

One important companion to this book is your own care team. Food and lifestyle education can support medical care; it cannot determine your diagnosis, medication doses, or personal nutrition needs. If you use insulin or medicines that can cause low blood sugar, changes in eating and movement deserve a coordinated plan. Kidney disease, pregnancy, eating-disorder history, and other medical needs also call for individualized guidance. We will make space for those realities throughout.

You have a life beyond the work of managing health. I hope these pages help you feel more able to enjoy it.

For now, let your shoulders drop a little, if they would like to.

We can begin where you are.

I. The care you have already been giving

There is a particular kind of disappointment in trying hard and having the numbers come back higher anyway.

You remember the walks you took. The cereal you stopped buying. The labels you read while someone waited behind you in the grocery aisle. You remember choosing the meal that seemed sensible when another one sounded better. Then a lab result arrives, and suddenly all that effort feels invisible.

The number deserves attention. So does the person reading it.

Before we talk about what to change, I want to acknowledge that you may already have been changing things for years. You may know more about food than you give yourself credit for. You may also be carrying a collection of instructions that do not fit together: eat more whole grains, eat fewer carbohydrates, eat breakfast, skip breakfast, choose fruit, be careful with fruit. A person could become exhausted before reaching the produce department.

Confusion in that situation is understandable. It does not establish that you are careless, and a health condition is not a character reference.

Consider the difference between an instruction and usable support. An instruction says, “Eat better.” Usable support asks what you currently eat, what you can afford, what you enjoy, who shares your kitchen, and what tomorrow requires. It gives you an action you can picture taking. It leaves you with a little more confidence instead of a larger pile of rules.

That is the kind of support I want Whole GI to offer.

It begins by treating your existing habits as information. If breakfast is toast, that is our starting point. If lunch disappears because you care for someone else, that matters. If your most reliable meal comes from a freezer, we have a freezer to work with. These are the conditions of your life, and any useful framework has to meet them.

You may recognize an old reflex here: the urge to tell me the whole history so I will understand why you have not done better. You do not owe me that defense. You are allowed to describe what is difficult without putting yourself on trial.

For the next few days, try replacing one judging sentence with a descriptive one. “I am terrible at lunch” can become “I often leave lunch too late.” The second sentence gives us somewhere to go. A sandwich prepared before an appointment could help. A reminder could help. A convenient meal could help. There are solutions available once the problem is small enough to name.

The same approach applies to the foods themselves. A breakfast can be enjoyable and need more staying power. A meal can be thoughtfully chosen and still leave you hungry. A routine can have served you well in another stage of life and need adjusting now. None of that erases the care behind it.

You get to keep the credit for the effort. You also get to ask for better tools.

Giving yourself a descriptive starting point has a practical benefit: you can choose a response that fits. You may notice less hesitation about a meal and more willingness to make one adjustment. That growing sense of capability is part of what you are building here. A routine becomes easier to return to when returning does not require an argument with yourself.

A small pause: Think of one thing you already do to care for yourself. Something ordinary counts. Before deciding what comes next, let that effort be visible.

2. What do you want your health for?

“I want my A1C lower” is a clear answer. “I would like my clothes to feel comfortable” is a clear answer, too. You do not need to apologize for either one.

But stay with the question for another moment. What would greater comfort, strength, or confidence make room for?

Maybe you want to take a trip without spending the weeks beforehand worrying about whether you will manage it. Maybe you want to carry groceries, wander a museum, return to tennis, work in the garden, or have something left in the tank after the errands. Maybe the answer is wonderfully small: you want to feel more like yourself at four in the afternoon.

That is the life inside the health goal.

We can become so accustomed to measuring the work that we forget why we began it. Numbers can give important information, yet they cannot describe the whole meaning of a day. A blood test cannot tell us how much you loved lunch with a friend. A scale does not know whether you felt confident enough to make a plan.

I like the phrase “the life you want to keep living” because it begins with something you already value. It assumes that your life contains people, pleasures, interests, and choices worth caring for. It does not require a glamorous answer. Enjoying your own home is an answer. Being able to ask for help and still direct your day is an answer. Making space for new things is an answer.

Independence deserves a generous definition here. Some people use mobility aids, rely on care, or live with limits that nutrition cannot remove. Your life remains yours. Agency can mean choosing a meal, deciding how help is offered, conserving energy for someone you love, or saying what matters to you. There is dignity in receiving support.

And your best healthy self does not have to be a younger self. You need not recover an old body to deserve a hopeful future. A useful goal can belong entirely to the person you are now.

Try finishing this sentence: “I want to care for my health so I can keep...” Give it a concrete ending. Keep walking to the ocean. Keep hosting Sunday lunch. Keep exploring somewhere new. Keep having enough attention for the conversations that matter.

You can use that answer when a food decision becomes noisy. Ask which choice supports the day you are trying to have. Sometimes that means cooking. Sometimes it means buying a practical shortcut and spending the saved energy elsewhere. Sometimes it means making a medical appointment, resting, or letting someone else take dinner.

Health care can be an act of participation in your own future. You are making room for the things you hope to do, while acknowledging that no food plan controls everything that happens to a body.

There is uncertainty in that. There is also purpose.

A goal connected to your life can help you decide which improvements matter most. Notice whether you are saving enough attention and effort for something you enjoy. You might begin planning an outing, making a meal you had stopped making, or asking for the support that would let you participate. Your health goals have a purpose beyond being monitored.

A small pause: Name one ordinary moment you would like more of in your everyday life. Let it be the reason behind one manageable act of care this week.

3. Blood sugar, without the blame

Let us give the science a chair at the table without allowing it to take over dinner.

Glucose is a fuel your body uses. Insulin, a hormone made by the pancreas, helps glucose move from the blood into cells. With insulin resistance, cells respond less effectively to insulin. The pancreas may compensate by making more; blood glucose can rise when that compensation is no longer enough. This picture involves biology, family history, activity, and other factors. It is much bigger than a decision about dessert.¹

That matters because people often hear a diagnosis as an accusation. They begin reviewing every food choice as if they are searching for the moment they caused this. There is more useful work available: understanding what your clinician has found and choosing the next support you need.

An A1C test estimates average blood glucose over roughly three months. A fasting glucose test describes a different piece of the picture. A1C is useful, but circumstances affecting red blood cells or hemoglobin can complicate its interpretation. When a result seems inconsistent with other readings, bring that question to your clinician.²

You do not need to become fluent in every laboratory abbreviation. It helps to know what is being measured, what target is appropriate for you, when the result will be reviewed, and what you should do between appointments. Write those questions down if remembering them in the room is difficult. Medical offices have a way of making very intelligent people forget every sentence they practiced in the car.

Food is one influence you can work with. So are movement, sleep, medication when needed, and the practical conditions that shape your routine. In the chapters ahead, we will keep returning to the meal because it gives us a concrete place to begin. We will also keep looking beyond it when the meal does not explain everything.

Please keep one distinction close: feeling tired, hungry, or foggy does not tell you your exact glucose level. Those experiences can have many causes. If monitoring has been recommended, use the method and targets your care team provides. A symptom is a reason to pay attention, not a reliable substitute for measurement.

The goal of learning is to make your questions better and your actions more useful. You might say, “My afternoon hunger is stronger on the days I miss lunch.” Or, “I have been waking repeatedly; could we discuss that alongside my glucose?” You are gathering context, rather than assigning fault.

And if medication is part of your plan, it belongs in this conversation without embarrassment. Good care can include several kinds of help. There is no prize for needing the fewest supports.

Some changes are felt, and others are measured. Less urgent hunger or a more comfortable afternoon can tell you that a routine feels better. Your glucose record and follow-up tests show how blood sugar is responding. Bring both to the conversation: what the numbers are doing, and what daily life feels like. You deserve care that pays attention to both.

For your next appointment: Ask, “What would you most like me to focus on before we meet again?” Then ask what that looks like in a normal day. A clear next step is worth taking home.

4. How a food ranking became a kitchen question

In 1981, David Jenkins and colleagues published work introducing the glycemic index as a way to compare the blood-glucose response to carbohydrate foods.³ The idea helped challenge a familiar assumption: foods with similar amounts of carbohydrate need not produce the same response.

GI compares the glucose response to a standardized amount of available carbohydrate in a food with a reference, commonly glucose. It reflects the measured response over time, not simply whether something tastes sweet. On the glucose scale, a GI of 55 or below is generally called low; 56-69 is medium; 70 or above is high. Values can differ with the food variety, processing, and preparation.⁴

This gives us a useful lens. It does not give us a complete lunch.

A ranking cannot tell us how much food you actually put on the plate, whether the meal supplies enough nourishment, or whether you can make it after an exhausting morning. It cannot know that you prefer rice, dislike a certain texture, share groceries with three people, or have ten minutes before leaving the house.

Glycemic load adds portion context. In its usual calculation, GL equals GI multiplied by the grams of available carbohydrate in the serving, divided by 100. Imagine a hypothetical food with a GI of 60 and a serving containing 15 grams of available carbohydrate. Its estimated GL would be 9. A serving with twice the carbohydrate would have an estimated GL of 18.⁴

That is a teaching example, not a forecast of your meter reading. It shows why the amount matters alongside the food's tested GI. You do not need to do arithmetic at every meal to understand the principle.

The next question is what happens when foods become a meal. We eat bread with things, rice with things, potatoes with things. We prepare them differently and eat them on different days, with different activity and medication patterns. A meal's practical effect cannot be captured by giving each ingredient a moral grade.

Whole GI grew from that kitchen question: how can a person use these ideas in a day they can actually repeat? The framework brings the portion, combination, preparation, and daily rhythm into view. It also makes room for the energy and appetite of the person doing the eating.

Whole GI brings that science into a framework I have spent thirteen years applying and refining with clients. Its principles have a purpose: build satisfying meals, manage the glycemic demands of those meals, support muscle, and give the day a rhythm you can maintain. The order and placement are deliberate. They connect what the research teaches us with what people need in order to use it.

You can learn a principle without turning it into another rule that follows you around. If you remember only one thing from this chapter, let it be this: a food makes more sense when you look at the serving and the meal around it.

The benefit of understanding these two ideas is freedom with direction. You can stop treating every carbohydrate as interchangeable and start making decisions about the actual meal in front of you. Notice when your questions become more specific: how much, with what, and where in my day? Those are the questions of someone learning how to make food work.

A small pause: Think about a carbohydrate you have been avoiding. Before deciding its place, consider the amount you enjoy, what you could serve with it, and any individual guidance you have received.

5. What the evidence can offer

There is a reason for the way the Whole GI day is built.

Protein at breakfast has a job. The carbohydrate portion and the foods around it have jobs. Giving daytime meals proper attention and making dinner more measured have jobs. So do movement, sleep, and the practical support that helps you keep feeding yourself well. These choices work together, rather than becoming a hundred separate health assignments.

A 2021 systematic review and meta-analysis of randomized trials in people with diabetes found that lower-GI or lower-GL dietary patterns improved glycemic control and several cardiometabolic risk factors. The average A1C difference was about 0.31 percentage points, with benefits also seen in measures such as LDL cholesterol and body weight.⁵ That is meaningful evidence for paying attention to carbohydrate quality and load within an overall eating pattern.

Whole GI combines this research with findings on protein, muscle, meal timing, food sequence, and the influences beyond food. Over thirteen years of working with clients, I have refined how those ideas fit into a usable day. Scientific evidence gives us a reason to choose a practice; experience with clients helps us teach it in a way that survives a busy kitchen, a small appetite, or a change of plans.

This is what I mean by a research-backed, client-tested framework. You are learning a considered set of practices, with a reason behind each one and room to adapt them to your own needs.

The holistic part matters. A more satisfying breakfast can make the morning easier. Adequate daytime meals can reduce the practical pressure to make dinner do all the feeding. An evening routine can create space for rest. Protein supports the muscles you use to move through your day. You are supporting several parts of health through a few connected decisions.

Nutrition guidance for diabetes emphasizes individualization: there is no single ideal balance of carbohydrate, fat, and protein for everyone.⁶ Whole GI gives that flexibility a structure. You have a dependable starting point, and you learn how to make it your own.

As you begin, notice what changes. Is hunger less urgent? Do meals leave you comfortably satisfied? Are cravings less insistent? Is your digestion more comfortable? How are you sleeping? Do you feel more able to prepare food and participate in your day? These experiences give you useful feedback about how the routine is serving you.

Keep that feedback alongside the measurements that describe your health. Feeling better is valuable in its own right; blood tests and glucose readings answer questions that feelings alone cannot. A more comfortable afternoon is encouraging, while a glucose trend needs appropriate measurement. Both kinds of information belong in the picture.

You do not have to notice everything at once. Nor must every useful change announce itself dramatically. Some benefits are quiet: breakfast keeps you satisfied, lunch happens before you are ravenous, dinner feels comfortable, and you spend less time wondering what you are allowed to eat. Let yourself recognize those things.

Keep close: Learn the reason, practice the principle, and notice the response. You are building understanding and confidence together.

6. Keep the foods you love

There may be a food you thought of as soon as you saw this chapter title.

Bread, perhaps. Rice. The pasta your family makes. The potato you like with butter and cinnamon. A particular breakfast that feels like home. Food carries memory, comfort, culture, and pleasure, as well as nutrients. A framework that ignores those things asks you to leave a large part of yourself outside the kitchen.

Here, we start by letting the food stay in the conversation.

Keeping a food does not mean that quantity, frequency, or medical needs disappear. It means we can examine its place thoughtfully before reaching for elimination. Sometimes the portion changes. Sometimes the meal needs a protein source or vegetables. Sometimes you discover a preparation you enjoy just as much. Sometimes a food remains an occasional pleasure, chosen with awareness rather than surrounded by guilt.

Start with the meal's identity. If you love oatmeal, what makes it oatmeal to you? Warmth, texture, cinnamon, the familiar bowl? Could you keep those qualities and include a protein source alongside it? If you love a sandwich, could the filling become more substantial and the vegetables more appealing? If rice belongs in your family's cooking, how could the rice fit into a meal with protein and vegetables rather than becoming a forbidden object?

Notice how the questions change the feeling. You are making something work. You are using knowledge to preserve enjoyment.

One practical exercise is to choose three meals you would be sorry to lose. Write down what you love about each, what feels difficult about fitting it into your health goals, and one adjustment you would consider. Keep the adjustment small enough that you would still recognize dinner.

A useful adaptation of spaghetti might retain the pasta, sauce, and shared table while changing the balance on the plate. A useful potato lunch might include a portion of potato with a satisfying protein topping and vegetables. A useful taco meal might include beans or another protein, vegetables, and an amount of tortilla that fits your plan. They keep the food recognizable while making the whole meal more supportive.

There is room here for pleasure that does not need an improvement project. You may enjoy a birthday dessert because it is a birthday dessert. A caring pattern is built across meals and days. One moment does not have to carry your entire health history.

When you want examples, the **Well + Easy recipe collection** is a place to browse familiar food. Choose one recipe that resembles something you already enjoy. You need not reinvent your kitchen to take the next step.

One early change to look for is that a favorite food stops creating such a long internal debate. You begin knowing how you want to serve it. You enjoy the meal and move on with your day. That confidence matters because a way of eating is easier to keep using when enjoyment remains part of it.

A small pause: “I would like to keep...” Complete that sentence with a food, a tradition, or a meal. Begin the adaptation there.

7. Look at the whole meal

Imagine looking into your refrigerator at lunch and finding a bowl of cooked rice. The question “Is rice good or bad?” can keep you standing there for a remarkably long time.

Try a more useful question: “What would make this a meal?”

Perhaps you have salmon, tofu, chicken, or beans. Perhaps there are frozen vegetables, leftover greens, or a cucumber waiting for a purpose. Perhaps olive oil, avocado, nuts, or a dressing would make the meal more satisfying. You are beginning to see components rather than a verdict.

For everyday planning, I often look for a recognizable protein source, vegetables or another fiber-containing food, an intentional carbohydrate portion, and some unsaturated fat. The exact balance depends on the meal and the person. Beans can contribute both protein and carbohydrate; nuts bring fat and some protein. Foods are wonderfully unwilling to stay in one tidy category.

This way of looking at a plate gives each food a purpose. Protein contributes to nourishment and fullness; vegetables, legumes, and other fiber-containing foods add useful substance; the carbohydrate portion supplies fuel; and healthy fat contributes flavor and satisfaction. The result is a meal designed to support you beyond the moment you finish eating.

Consider three ordinary upgrades. A salad with leaves and a little dressing could become lunch with beans, fish, eggs, chicken, or tofu and an appropriate carbohydrate choice. Toast and coffee could become a more substantial breakfast with a protein food and produce. A bowl of vegetables at dinner could become adequate nourishment when you add the protein and other food your day requires.

You may notice that some of these changes add food. That can feel unfamiliar if caring for weight has always meant removing things. The

question is whether the meal meets your needs. A sparse plate can look virtuous while leaving a person poorly fed and thinking about snacks all afternoon.

Fiber deserves a place in this picture. A systematic review in people with diabetes found benefits of higher-fiber diets for several measures of metabolic health.⁷ That supports making room for foods such as legumes, vegetables, fruit, and whole grains where they suit you. Increase fiber gradually if needed, and follow any condition-specific guidance when higher fiber is not appropriate.

Healthy fat has a place for more than flavor. The PREDIMED trial found fewer major cardiovascular events among high-risk adults assigned to Mediterranean diets enriched with extra-virgin olive oil or nuts than among those assigned to a reduced-fat comparison diet.³¹ This supports the choice of foods such as olive oil and nuts within a nourishing pattern. Some benefits show up in long-term health measures rather than an immediate feeling after lunch.

For now, practice noticing what is present and what may be missing. You can do that without weighing every ingredient. If you need carbohydrate counting or other detailed tracking for medical care, keep using it; this broader view can sit alongside it.

Over the next week, notice how long the meal feels satisfying and how hunger returns. Does a more substantial salad serve you better than leaves alone? Does a breakfast with an obvious protein source carry you more comfortably toward lunch? You are learning the difference between having eaten something and having fed yourself well.

Fiber and fluids also work together to support bowel regularity.²⁵ As you introduce fiber gradually, notice whether stools become easier to pass and your routine more comfortable. Persistent discomfort is a reason to adjust the approach and ask for help, rather than pushing through a larger and larger bowl of bran.

At your next meal: Identify the protein first. Then look for produce or another fiber source, your carbohydrate portion, and the rest of the meal. Choose one useful adjustment rather than correcting everything at once.

8. Preparation, portions, and a little perspective

The same ingredient can arrive in very different forms. A grain may be intact, milled, puffed, or cooked until soft. A potato may be part of a mixed meal or a large serving eaten on its own. This is one reason a food's name does not tell the entire story.

Tested values can differ across products.⁴ A reference is most useful when the tested food closely resembles what you eat.

You do not need to memorize the tables. You can develop a few dependable choices that you enjoy and understand. A bread you like, a grain that cooks reliably, a bean you will actually use, a breakfast that fits your morning. Familiarity reduces the number of decisions a meal requires.

Preparation is part of this learning, but it deserves perspective. You may hear about cooling starches or changing cooking methods to alter their effect. These ideas are useful refinements within the larger picture of portion and meal composition. You do not need to perform a special ritual before enjoying lunch.

Preparation can change how some starch is digested. In a small crossover study, cooked white rice that was refrigerated for 24 hours and reheated contained more resistant starch and produced a lower glycemic response than freshly cooked rice.³² This is one practical reason preparation belongs in Whole GI. You can use safely stored leftovers as part of an intentional meal, with the portion and its companions still in view. Follow the cooling and reheating guidance in Chapter 21.

Meal sequence is another idea worth discussing carefully. Clinical meal studies have found that beginning with protein and vegetables before the carbohydrate portion can reduce the immediate glucose response.⁸

When the meal naturally allows it, this is a simple way to put the same food to work differently. You can begin with the vegetables on your plate without dismantling a sandwich or policing every bite.

If you like beginning a meal with vegetables or protein, you can do that. If you are eating a mixed dish, enjoy your mixed dish. The aim is a routine you can carry into an ordinary life without becoming anxious about the order of your forkfuls.

Portions are easier to discuss when we remove the courtroom atmosphere. A serving is information. It helps you learn what feels satisfying, what fits your individual plan, and what the rest of the meal needs. You can use a familiar bowl or serving spoon as a practical reference. You can also use more precise measures when your medical care calls for them.

Try changing one variable at a time when you are learning. If you change the food, the portion, the timing, and the preparation together, it becomes hard to know which change helped or which one made the meal less enjoyable. Give yourself a chance to learn something specific.

The benefit of meal sequence is that it can change the immediate glucose response without requiring a completely different meal.⁸ If glucose monitoring is part of your care, comparable meals can give you something specific to discuss with your clinician. You may feel no obvious difference while a measured response changes. A useful practice does not have to produce a sensation to be doing something worthwhile.

A small pause: Choose one familiar meal to practice with this week. Keep most of it the same. Make one adjustment you can describe in a sentence.

9. A rhythm for an ordinary day

A framework is most useful when you can picture it before you need it.

The Whole GI day begins with hydration, moves into a protein-forward breakfast, makes lunch a substantial meal with a place for starchier carbohydrates, and uses a dinner centered on protein, healthy fat, and non-starchy vegetables. Snacks and dessert are optional. Herbal tea can provide a pleasant close to the day.

These are landmarks. They give you a starting shape, so every meal does not require a fresh negotiation.

The fuller program teaches a breakfast guideline of at least 25 grams of protein within about 60-90 minutes of waking, and emphasizes lunch as the largest meal. They provide a clear starting point, with timing and amounts adapted to your appetite, schedule, activity, and care needs.

Meal timing matters because your body has a daily metabolic rhythm. In a controlled study in men with prediabetes, an earlier eating schedule improved insulin sensitivity even without weight loss.⁹ Another randomized trial in adults with type 2 diabetes found lower daily hyperglycemia when more food was placed at breakfast and less at dinner.²³ These findings support the Whole GI emphasis on feeding yourself properly earlier in the day and easing the load at night. The framework translates that direction into ordinary meals.

What we can do is look at your day. Do you leave so much food until evening that dinner becomes the first real meal? Does breakfast disappear into coffee? Is lunch too small to support the afternoon? Does a snack help, or is it merely an item on a schedule? These questions make timing practical.

A useful daily rhythm should also be humane. It should accommodate the person who rises early and the person who works nights. It should leave

room for appetite and for eating enough. The framework can be adapted with professional help when a standard pattern does not fit.

The purpose of each part of your day

Morning hydration begins the day with a basic need attended to. A randomized trial in young adults found that rehydration after water restriction improved aspects of mood and cognitive performance.²⁹ A glass of water on waking is an easy cue to begin meeting your fluid needs. Notice thirst, comfort, and whether drinking regularly becomes easier. Use your prescribed fluid plan when you have one.

A protein-forward breakfast is there to support nourishment and satisfaction early in the day. Breakfast research shows benefits for fullness and appetite control.²⁰²¹ Look for a morning that feels less driven by cravings and hunger, with enough room in your attention for something beyond the next snack.

The optional midmorning snack bridges a gap when needed. Research supports attention to snack composition and appetite.³⁵ Notice whether it leaves you comfortably ready for lunch, with its amount and timing serving your day.

Lunch brings protein, vegetables, healthy fat, and an intentional starch portion together in a substantial daytime meal. It helps you use carbohydrate quality and load within the pattern supported by low-GI/GL research.⁵ Notice satisfaction and the ease of moving through the afternoon. Chapter 11 helps you build it from food you recognize.

The optional afternoon snack helps when there is a long interval before dinner or a small appetite makes another eating occasion useful. In a crossover study, a high-protein yogurt snack improved appetite measures compared with common higher-fat snack foods.³⁵ Look for less urgent hunger while dinner is being prepared, rather than an extra eating occasion you feel obliged to fit in.

Dinner brings protein, healthy fat, and non-starchy vegetables into the evening, with a more measured carbohydrate load. Clinical meal-timing

research gives a reason to avoid placing the day's heaviest eating close to bedtime.²²²³ Notice comfortable satisfaction, evening appetite, and sleep. Chapter 12 connects those observations to the whole evening routine.

An optional evening snack or dessert gives pleasure and nourishment an intentional place. Portion and the amount of carbohydrate remain part of the meal picture, consistent with individualized diabetes nutrition guidance.⁶ Notice whether your choice satisfies you and lets the eating occasion feel complete. If you need a bedtime snack for medication-related reasons, its role comes from your care plan.

Herbal tea can close the day with a caffeine-free drink and a familiar pause. A randomized study found that a substantial caffeine dose could disturb sleep even when taken six hours before bed.³⁰ Choosing a suitable caffeine-free evening drink protects that part of the routine. Notice whether it helps you settle into a quieter evening; choose herbs that are compatible with your medicines.¹⁹

These parts support one another. You are feeding the morning, making room for lunch, easing the evening, and giving recovery a place. The benefit is a connected day, with fewer last-minute decisions and several health priorities supported by the same ordinary meals.

Here is a gentle place to begin: make your next main meal easier to anticipate. Know what it will be, where the protein will come from, and whether you have the ingredients. Once that becomes familiar, look at the meal before it or after it. A day takes shape through small connections.

You can explore the [Daily Framework](#) when you want the morning-to-night overview. The [Meal Planner](#) gives those parts of the day a place to live. You do not need to fill every space to have used the tool well.

The aim of the daily flow is to support nourishment, appetite, and the demands of the body across the day. In practice, notice whether your meals begin to feel connected. Is breakfast helping you reach lunch comfortably? Is a proper lunch making dinner feel less urgent? Are you spending less of the evening trying to catch up on the food you missed earlier? These are useful signs that the rhythm fits your life.

A small pause: Which part of your day would be most relieving to make predictable? Start there. A reliable lunch is a perfectly good doorway into Whole GI.

10. Breakfast you can recognize

Morning is a poor time to ask a tired person to invent a new identity.

If you like a warm breakfast, let us work with warmth. If you prefer something cold, assembled, savory, or taken with you, those preferences belong in the plan. A breakfast you dislike is unlikely to become a reassuring daily ritual, however impressive it looks in a photograph.

The Whole GI approach begins by giving protein a visible role, alongside a suitable carbohydrate and healthy fat. You might use eggs with another protein food, plain Greek yogurt, cottage cheese, tofu, or another option that fits your preferences. Amounts and product labels matter; a spoonful of nuts or a splash of plant milk may contribute less protein than you expect.

You do not have to calculate forever. You can check a few dependable combinations once and become familiar with them. Some people enjoy the precision of numbers; others find it relieving to know their usual breakfast is well considered. Choose the level of detail that helps you, while retaining any tracking needed for your care.

For an oatmeal morning, you might keep the oats and cinnamon and include a substantial protein food alongside them. For toast, you might add eggs, cottage cheese, or another protein filling and some produce. For a yogurt bowl, choose a product with useful protein, then consider fruit and nuts or seeds in portions that suit you. Plant-based eaters can build around tofu or other protein-rich options, checking the label when products vary.

There is no need to stage the bowl. You can eat it from the same slightly chipped favorite you have used for years. It has done good work.

If your appetite is small, a full breakfast may feel like a challenge. A smaller, nourishing portion or a planned second eating occasion can be a more workable beginning. Medication schedules and conditions affecting

digestion deserve specific guidance. We will return to small appetite later in the book.

One useful household decision is to choose two breakfasts: a usual one and an easy backup. The backup should be possible on a morning when the kitchen is busy and you cannot find your glasses, although they are, naturally, on your head. Keep its ingredients available. Let repetition do some of the work.

The planner and recipes can help you find combinations, but the result belongs to you. If your reliable breakfast comes from four familiar ingredients, it is doing exactly what we hoped: making a useful choice feel ordinary.

Protein has a deliberate place at the beginning of the day because fullness is one of its useful effects. In a randomized crossover study of young adult women, a protein-rich breakfast increased fullness and satisfaction and reduced hunger during the hours before lunch.²⁰ Another clinical study, in breakfast-skipping adolescent girls, found improvements in appetite control and reduced evening snacking with a higher-protein breakfast.²¹ The research supports making breakfast more substantial rather than relying on willpower later.

You may notice that food becomes less insistent in your thoughts. The midmorning search for something sweet may ease. You may reach lunch hungry enough to enjoy it without feeling that you need food immediately. These are the kinds of changes to look for when you give breakfast enough protein and enough overall nourishment.

Keep a simple comparison in mind: how does your usual morning feel, and how does it feel after a week of a more complete breakfast? Notice fullness, cravings, satisfaction, and the ease of getting on with what you wanted to do. You do not need an appetite scorecard attached to the refrigerator. A short observation is enough.

Tomorrow morning: Decide tonight what breakfast will be. Put one ingredient, bowl, or reminder where you will see it. Make the next step kind to your waking self.

11. Give lunch a proper place

Lunch often receives whatever attention is left over.

It gets squeezed between appointments, assembled from bits, or postponed until it begins competing with dinner. People who would never forget to feed a child, a guest, or a pet can become surprisingly negotiable about feeding themselves.

Your lunch deserves a place in the day.

Within Whole GI, lunch is deliberately substantial. It is where I teach a clear place for starchier carbohydrates alongside protein, vegetables, and healthy fat. That might look like a sandwich, a bowl with rice, a potato with a nourishing topping, a bean-based meal, or pasta with a more balanced set of companions.

This is a useful place to practice keeping the foods you love. You can take the familiar starch out of the category of “something I should probably avoid” and begin learning its portion and combination. Medical guidance still takes priority, and the meal should suit your own needs.

Consider a bowl assembled from leftover chicken or tofu, vegetables, a serving of grain, and a dressing you enjoy. Consider a sandwich with a substantial filling and a side of produce. Consider bean soup with an additional protein food if needed and a carbohydrate portion that fits your plan. None requires a separate cuisine called healthy food.

The important logistical question is when lunch becomes real. If you wait until you are hungry to discover there is nothing prepared, you have created an unnecessarily difficult decision. A little planning can happen earlier: set aside dinner ingredients for tomorrow, choose a dependable shop-bought option, or keep the components of an assembled meal ready.

For someone working outside the home, lunch may require a container and access to safe storage. For someone at home, it may require treating

the meal as an appointment worth keeping. For someone caregiving, it may require making your own plate when you make another person's. The barrier determines the solution.

A substantial lunch is not an instruction to eat beyond comfort. If you have a small appetite, need a different distribution, or follow a medical meal plan, adapt. The useful principle is to stop treating daytime nourishment as optional background work.

Try building lunch around something you already have. When you choose recipes in the **Meal Planner**, look for overlap in ingredients and opportunities to use leftovers. Even two connected meals can make the week feel less fragmented.

Lunch also gives you another opportunity to nourish your muscles rather than leaving most protein until dinner. Research in community-dwelling adults aged 55 and older found that more protein at breakfast and lunch was associated with higher total daily protein intake.²⁷ This is one reason a substantial daytime meal deserves attention as we age.

As lunch becomes more dependable, look for less urgent afternoon hunger, less need to graze while preparing dinner, and a more comfortable evening appetite. Those are practical observations to make. If lunch consistently leaves you sleepy or uncomfortable, review its size, composition, and the rest of your day. The aim is a meal that supports the afternoon you want to have.

A small pause: What usually interrupts lunch? Name the interruption before choosing the recipe. The best meal idea is one you can actually reach.

12. Dinner can be simple and still be dinner

By evening, a recipe with twelve separate chopping instructions can feel personally insensitive.

Dinner needs to respect the energy available at the end of the day. Within the usual Whole GI flow, it centers on protein, healthy fat, and non-starchy vegetables, with less emphasis on the starchier carbohydrate portion used at lunch. It should still be enough food.

Dinner should leave you comfortably nourished. The aim is a satisfying evening meal that complements what you have eaten during the day. Enough protein, vegetables you enjoy, and useful flavor make that possible without asking dinner to become your largest eating occasion.

Think of an evening meal in components: a protein you can prepare or assemble, vegetables in a form you enjoy, and flavor. Fish with frozen green beans and olive oil. Tofu with a bag of stir-fry vegetables and a sauce that suits you. Chicken with roasted vegetables. An egg-based meal with a vegetable side. A plant-based dinner may include legumes and therefore carbohydrate; food categories should help us think, rather than stop us from eating sensibly.

You can share a dinner even when plates differ. One person may want an additional starch portion; another may need a different amount. Put components on the table and let the meal support the people present. You do not need two complete cooking operations.

If dinner is also your favorite social meal, keep the social part. Sit down when possible. Use a plate you like. Let conversation be allowed to outrank nutritional commentary. No one needs a running analysis of the broccoli.

On an easier evening, make an extra portion of a component for tomorrow. On a difficult evening, take the shortcut. A microwave, a bag

of frozen vegetables, or a prepared protein can be part of a cared-for meal. We are trying to feed a person, and that includes the person doing the preparation.

It may help to keep a short list of dinners that require very little thought. Three is enough to begin. One you cook, one you assemble, and one you buy or bring home. Know what you like from each option before an exhausted evening asks you to decide.

The evening composition has a scientific reason. A dinner built around protein, healthy fat, and non-starchy vegetables can reduce the carbohydrate load compared with a starch-heavy dinner while still supplying useful nourishment. Timing matters alongside composition: a randomized crossover trial found higher overnight glucose after a late dinner than after the same dinner eaten earlier.²² Whole GI brings those considerations together so the evening meal supports the night's work of rest and recovery.

Sleep is worth paying attention to here. In my work with clients, better sleep is one of the changes reported as the Whole GI evening pattern becomes established. Notice whether you settle more comfortably, wake less often, or feel more rested in the morning. Consider the whole evening: how much you ate, when you finished, whether you felt comfortably satisfied, and what you drank. Research also connects carbohydrate intake with aspects of sleep.²⁴ Together with what clients report, that makes your nights a useful part of the picture to observe.

Comfort also matters. For people with nighttime reflux, leaving at least three hours between a meal and lying down may improve symptoms.²⁶ A comfortable evening meal with room before bed can therefore support a different part of the night as well. One ordinary dinner can be part of caring for several things at once.

Look for a meal that leaves you satisfied without feeling heavily full, an evening with less unplanned eating, and nights that feel easier. If you use glucose monitoring, bring the overnight and morning patterns to your care team as well. These are related pieces of feedback, each telling you something useful.

Tonight: Ask, “What is the simplest dinner that will meet my needs?” Let the answer be simple without making it inadequate.

13. Snacks, dessert, and the end of earning food

Somewhere along the way, many of us learned to describe eating as a series of moral events. We were good at breakfast, bad at lunch, and considering whether dinner could repair our reputation.

It is a tiring way to spend a day.

Snacks can have a purpose. They may bridge a long gap, respond to hunger, help you eat enough when appetite is small, or fit a medication plan. They do not need to appear simply because a schedule has a space for them. In Whole GI, they are optional unless your individual care plan says otherwise.

A practical snack might combine a protein food with another useful component: plain yogurt and berries, an egg with vegetables, cheese with fruit, or another combination that fits your needs. Choose a portion that serves its purpose. If afternoon hunger is reliably intense, also look back at lunch. A snack can help today while the larger pattern deserves attention.

Dessert asks for a similar kind of honesty. Do you want it? Will it be enjoyable? What amount feels appropriate? What does your care plan require? You can answer those questions without naming the food a cheat.

The framework makes room for an intentional dessert rather than treating sweetness as automatic failure. Having it with or after a meal can help you make it an intentional part of eating, with the portion considered alongside the rest of the meal. A smaller portion of something you really enjoy can be more satisfying than several substitutes you never wanted.

That is also permission to stop when something is disappointing. You do not have to finish a joyless dessert in the name of discipline. The freezer can keep its chalky mystery square.

There is one medical distinction to keep clear. A low-glucose episode is treated with the fast-acting carbohydrate and response plan recommended by your care team. A balanced snack, a low-GI dessert, or a protein food is not interchangeable with that treatment. If lows occur, ask for a review of the plan.¹⁰

When the day closes, a cup of herbal tea can be a pleasant ritual if you enjoy it. The familiar sequence of making it, sitting down, and letting the evening quieten can give rest a place in the day. Herbs can interact with medicines, so choose with care and ask a pharmacist about products you use regularly.¹⁹ Plain water is welcome, too. Choose water in a form you enjoy and follow any prescribed fluid guidance.

Snack composition has been examined directly in clinical studies. In a randomized crossover study of healthy women, a protein-rich yogurt snack improved hunger or fullness measures compared with common higher-fat snacks.³⁵ The practical lesson is to make the snack useful: choose nourishment and an amount that serves the interval, rather than relying on grazing to carry you through the afternoon.

For dessert, the principle is thoughtful inclusion. An enjoyable portion, considered within the meal's carbohydrate amount, can keep pleasure in a routine you want to continue.⁶ Pay attention to satisfaction and any measured glucose patterns relevant to your care. A food can have a place without becoming an all-or-nothing decision.

For the evening drink, protecting sleep begins partly with what you leave out of the cup. Clinical research on caffeine supports choosing a caffeine-free option later in the day.³⁰ Your herbal tea is both a practical drink choice and a cue that the day's work can begin to end. You do not need a new project at bedtime.

The benefit of an intentional snack is that it answers a need before the need becomes overwhelming. Notice whether it comfortably bridges the gap, helps you arrive at the next meal with a manageable appetite, or makes it easier to eat enough on a small-appetite day. With dessert, a useful change may be satisfaction: you choose something you enjoy, have an amount that fits, and feel able to let the meal end.

A small pause: Notice one food word that makes you feel judged. Could you replace it with a useful description: satisfying, convenient, sweet, substantial, occasional, or simply enjoyable?

14. When today has less to give

There are days when the recipe is not the problem. Standing at the counter is the problem. Opening the container is the problem. Driving to the store is the problem. Remembering that the chicken is still frozen is an additional problem, although the chicken seems untroubled.

A food plan should recognize the work required to use it.

Cooking time is only one kind of effort. Chopping, lifting, reaching, washing dishes, and making decisions all draw on something. A meal that takes fifteen minutes can still be difficult for a person with painful hands or limited stamina. Your circumstances belong in the plan before the recipe is selected.

Begin by asking what kind of effort you can manage today. Can you assemble a meal? Use the microwave? Sit for preparation? Reheat something already cooked? Would a ready-to-eat option or help with delivery solve the most important barrier?

Then choose accordingly. Prepared chicken or tofu, canned fish, cooked eggs, frozen vegetables, washed greens, canned beans, and other convenient ingredients may give you useful components with less work. Check labels for your particular nutrition needs. Choose packaging you can open and portions you can store. Convenience becomes valuable when it makes nourishment reachable.

It helps to give your low-energy meals the dignity of a plan. Choose three that you enjoy while you have the attention to do so. Keep the ingredients available when possible. Write the combinations somewhere easy to find. The list is there to spare you another decision on the days when decisions are expensive.

For example, your assembled lunch might use canned salmon, a suitable ready-cooked grain portion, and a vegetable you can manage. Your

dinner might be a prepared protein with microwave vegetables and a familiar dressing. Your small breakfast might use a protein-rich yogurt and another component that suits you. Adapt the foods to allergies, medical needs, and taste; the practical lesson is that a meal can be assembled.

You can also make the environment more helpful. Move the foods you use often within reach. Keep a stable seat available for appropriate prep tasks. Use safe tools that suit your hands. Ask someone to help with the physically difficult part rather than vaguely asking them to “help more.” Specific requests are easier to fulfill: open these containers, portion that food, carry this bag, wash the pan.

Receiving help does not make the meal less yours. It can preserve your ability to decide what you eat and how you spend the rest of your energy.

The [Special Diets collection](#) and the planner's preferences can help you narrow choices when simpler preparation or smaller meals would help. Choose a few useful filters; a long list of restrictions can hide meals you could easily adapt.

This is the day the framework needs to fit. You do not have to wait for a better version of yourself to deserve a proper lunch.

On these days, success can look like getting adequately fed without using all the effort available to you. Notice fewer missed meals, less time standing in discomfort, or more attention left for the rest of your day. Those are practical benefits of adaptation. Easier food can make the difference between knowing what nourishment matters and being able to receive it.

Make a rescue list: One breakfast, one lunch, one dinner. Each should be possible with the effort you can manage on a difficult day. Keep the list short enough to use.

15. A small appetite still deserves nourishment

“I am just not very hungry” can sound like a convenient development in a world that talks constantly about eating less.

It can also make feeding yourself surprisingly difficult.

When appetite is small, the challenge may be getting enough food, protein, and other nutrients into the amount you can comfortably eat. Larger salads, bigger glasses of water before meals, and instructions to fill up may be poorly matched to the situation. Less appetite does not automatically mean fewer nutrition needs.

Clinical nutrition guidance for older adults emphasizes adequate nourishment and hydration, attention to malnutrition risk, and individualized support.¹¹ If appetite has changed persistently, weight is falling unintentionally, or eating is becoming difficult, bring it to your clinician. The cause deserves attention; you should not have to solve it through recipes alone.

On the practical side, begin with a smaller meal that includes a useful protein source. Choose textures, temperatures, and flavors that appeal to you. A modest bowl may feel more inviting than a large plate. A smaller meal now and another planned eating occasion later may be easier than trying to finish everything at once.

Consider compact combinations you already enjoy: a small egg dish with a suitable side, a protein-rich yogurt bowl, fish or tofu with a little grain and vegetables, or a nourishing soup paired with something that supplies what the soup lacks. A cup of food is not automatically complete; look at what is inside it.

If you use a GLP-1 medicine, the food conversation belongs alongside your prescribing care. A joint professional advisory highlights nutritional adequacy, attention to muscle and function, and supportive habits

during treatment.¹² This book cannot determine a dose, a target intake, or how to stop a medicine. It can help you describe what is happening: what you can eat, what feels difficult, and what support you need.

Do not force a large or rich meal through nausea in order to follow a general framework. Persistent vomiting, inability to keep fluids down, significant pain, or concerning weakness needs prompt medical advice. After stopping a medication, appetite and routines may change again; planning with your clinician gives you somewhere to bring those changes.

There can be an emotional adjustment, too. Someone who has spent years trying to eat less may find it strange to hear “you need enough.” Enough is a caring word. It makes room for your muscles, your attention, and the ordinary tasks you want to do.

The Well + Easy collections are a place to explore meals suited to different needs, including smaller-meal options. Use the choices that fit your needs rather than treating a label as a complete medical recommendation.

The purpose of a smaller, thoughtfully built meal is to make useful nourishment achievable within your appetite. Notice which portions you can comfortably finish and which combinations help you eat regularly without feeling overfaced. Over time, adequate intake and stable function matter more than the visual size of the plate. Bring ongoing appetite changes, weakness, or unintended weight loss to your care team so the support can be tailored.

A small pause: Is the difficulty quantity, appetite, chewing, nausea, fatigue, cost, or access? Naming the actual barrier helps you ask for the right kind of help.

16. Strength belongs in the conversation

When you picture getting stronger, you may picture a gym. You might also picture lifting a suitcase into a car, carrying a pot to the sink, or getting up from a low chair without needing quite so much planning.

Strength has ordinary jobs.

That is why I want it included whenever we talk about weight. A smaller number can be meaningful for some goals, but it cannot tell us whether you are adequately nourished or maintaining the ability to do the things that matter to you. The body we care for is a working body, even when its work looks quiet from the outside.

Protein and appropriately tailored exercise both matter for maintaining muscle function with age. Expert guidance emphasizes their combination and the need to account for health status.¹³ Protein needs differ, and kidney disease or other conditions can change recommendations. You do not need to adopt someone else's target from a video.

Practically, give protein a regular place in meals rather than leaving nearly all of it until dinner. Choose foods you like and can obtain. If animal foods are part of your routine, you have several options. If you eat plant-based meals, plan deliberately around useful protein sources rather than assuming any vegetable dish supplies enough.

Movement should also match your body. A qualified professional can help you find safe strength work, including adaptations for pain, balance, disability, or recovery. You do not have to prove you can perform a standard exercise before asking for a version you can use.

There is a difference between a challenge that helps and an instruction that ignores your limits. You are allowed to ask which one you are being offered. You are also allowed to value gradual progress, rest, and

rehabilitation. Confidence often grows through finding something you can repeat safely.

If weight loss is appropriate for you, discuss how to pursue it while protecting nourishment and function. If maintaining weight or regaining lost weight is more appropriate, that belongs here equally. The aim is care for your actual body, not loyalty to a goal that no longer fits.

You may find it useful to notice a functional change alongside the usual numbers. Is a familiar activity becoming easier? Are you better able to prepare lunch? Do you recover more comfortably after an outing? Those observations do not replace clinical assessment, but they can help describe what health means in your daily life.

Please resist turning every activity into a test. You can enjoy the walk without timing it. You can dance badly in the kitchen without collecting data. There is room for movement that feels like living.

Protein supplies the building material for muscle; appropriately challenging movement gives the body a reason to maintain and build it. A meta-analysis of randomized trials found that protein supplementation added to resistance training improved gains in muscle size and strength in healthy adults.²⁸ That helps explain why Whole GI keeps nourishment and movement in the same conversation.

Notice progress over weeks, through activities that matter to you. A familiar exercise may feel more manageable, or a task may require less effort. You might feel more confident getting up from a chair or carrying something that used to feel awkward. Choose observations appropriate to your abilities, and let a professional help you assess change when needed. Strength is often recognized in the middle of living.

A small pause: Name one task you would like to feel more capable doing. Ask what nutrition support, movement guidance, equipment, or practical help could make it more accessible.

17. A changing body deserves a changing plan

You may have noticed that the routines that once felt dependable no longer give you the same experience. Your shape may have changed. Sleep may have become less predictable. Your appetite, recovery, or energy may feel unfamiliar.

It is reasonable to want an explanation. It is also understandable to feel irritated by another recommendation to simply try harder.

For women in midlife, aging, the menopause transition, activity, sleep, and other factors can all be relevant to changes in weight and body composition. The Menopause Society describes this as a multifactorial picture, rather than a single food mistake.¹⁴ You deserve support that takes the whole picture seriously.

Whole GI does that through a daily food framework and a second layer of attention to sleep, stress, inflammation, and hormonal factors. The program calls its self-assessment the Root Cause Finder. It helps you identify patterns worth addressing and bring clearer questions to your care team. The assessment guides reflection and education; a clinician handles diagnosis.

This distinction protects you from a different kind of overwhelm. You should not have to conclude that your hormones, cortisol, and inflammation all need fixing before you can eat breakfast. You can feed yourself today while bringing persistent symptoms to a clinician who can investigate them.

Try describing the changes in ordinary terms. “I wake several times most nights.” “My appetite changed after this medication.” “I am struggling with hot flashes and fatigue.” “Pain has made shopping harder.” Specific observations give a care team more to work with than a broad conclusion that everything is out of balance.

They also help you choose practical support. If nights are difficult, an easy breakfast may matter more than an elaborate meal plan. If pain limits shopping, access deserves attention. If a medicine affects appetite, your food pattern may need adapting. The same symptom can have several explanations, so avoid diagnosing yourself through a checklist.

Men may encounter changes in appetite, activity, sleep, and health needs, too. The framework's practical questions still apply. Partners can learn together without assuming identical portions or medical goals. Sharing a table does not require sharing a prescription.

The hopeful part is that a changing body can invite a more appropriate plan. You can stop asking a routine from twenty years ago to solve every current problem. You can bring new information into the way you care for yourself.

The Whole GI program overview introduces the wider approach. Use it as a companion to your care, with room for professional assessment when a question needs more than education.

The benefit of this wider view is that you can stop asking food restriction to solve every difficulty. An adjustment to symptom care, a more workable meal pattern, or help with physical access may make several things easier together. Notice which change gives you more capacity: a better night that makes breakfast possible, less discomfort that makes shopping easier, or an eating routine that fits a changed appetite.

A small pause: What has changed? What support would make that change easier to live with? Let the second question be as important as the first.

18. Beyond the plate, without another project

There is a point at which health advice becomes a second occupation. You are supposed to plan meals, improve sleep, manage stress, move more, drink water, make appointments, and somehow remain wonderfully relaxed about the workload.

Let us approach the rest of the day with a little more kindness.

Sleep is one reason food may not explain the whole glucose picture. In a randomized trial, sustained modest sleep restriction impaired insulin sensitivity in women.¹⁵ That gives sleep a clear place in metabolic care, alongside the food on your plate.

If nights are difficult, choose a useful next question. Is the problem opportunity to sleep, discomfort, hot flashes, worry, frequent waking, or symptoms such as snoring and daytime sleepiness? A routine can help with some barriers; others need medical attention. You are not failing at sleep because a bath and a lavender pillow did not resolve them.

Stress deserves similar respect. Some of it is an appointment you can move. Some of it is caregiving, financial pressure, grief, illness, or work you cannot simply put down. The useful question is what support can reduce the load or help you get through it. A prepared lunch may be one form of support. So may a conversation, a boundary, a service, or help from another person.

Inflammation and hormonal changes are part of the wider health conversation, too. Whole GI helps you look for the useful connections: pain that limits movement, symptoms that interrupt sleep, or changes that make eating difficult. Addressing those patterns with the right professional can make your daily food routine easier to use.

Movement is another practical companion to meals. A randomized crossover study in adults with type 2 diabetes found that advice to walk

after meals improved post-meal glycemia more than advice that did not specify timing.¹⁶ That gives us a simple tool that can fit into an ordinary day. Choose safe movement within your abilities and discuss adaptations when needed.

If walking is difficult, ask what is appropriate for you. Brief activity breaks have also been studied in people with type 2 diabetes, including light walking and simple resistance activities.¹⁷ Choose an appropriate version with your clinician or movement professional when you need guidance.

You can begin with one area. Perhaps you arrange a sleep conversation, prepare lunch earlier, ask for a movement referral, or make the evening less crowded. Choose something that removes friction from the day you actually have.

The four areas beyond food

Sleep supports how the body responds to insulin, which is why it belongs inside the same framework as meals.¹⁵ A regular opportunity for rest, attention to symptoms that interrupt sleep, and an evening routine you can repeat give that principle somewhere to live. Notice how rested you feel on waking and whether you are waking less often. Those observations can also help your clinician identify what needs attention.

Stress and cortisol belong in the wider picture because the stress response and glucose regulation interact. A randomized study in adults with type 2 diabetes found that a structured stress-management program improved A1C over a year.³³ Whole GI makes room for the practical version: a pause, a manageable plan, support with an overwhelming task, and care for ongoing distress. Notice less tension around meals and a greater ability to respond deliberately. These everyday practices can accompany more structured support when that is useful.

Inflammation is one reason the quality of the whole eating pattern matters. A randomized trial in people with metabolic syndrome found

improvements in inflammatory markers with a Mediterranean-style diet.³⁴ The overlap with Whole GI is useful: vegetables, fiber-rich foods, and healthy fats serve more than one purpose. You do not need to feel inflammation changing in order for a nourishing pattern to be worthwhile; your clinician can interpret relevant health measures.

Hormonal changes, especially through menopause, can affect body composition and the circumstances of eating, sleeping, and moving.¹⁴ Supporting protein intake, appropriate strength work, sleep, and symptom care gives you a coordinated response.¹³ Notice what becomes easier when the plan fits your current body: more workable meals, a more manageable appetite, or greater capacity for movement and rest. Keep symptom treatment part of that conversation when needed.

The same few practices appear across these areas because the body is connected. You do not have to run a sleep program, a food program, a muscle program, and a stress program as four separate jobs. You are building a daily flow that gives each priority some support.

These connections are the heart of a holistic framework. Sleep belongs in the blood-sugar conversation because of its relationship with insulin sensitivity.¹⁵ Movement belongs because it helps muscles use glucose.¹⁶ Food belongs because it supplies nourishment and shapes the demands of a meal. You are supporting overlapping systems through a small set of repeatable practices.

Watch for patterns across several days: feeling more rested, hunger that is easier to respond to, a meal routine that takes less effort, or movement becoming more regular. If you monitor glucose, look at the trends with your clinician. A short walk and a simpler dinner can fit into the same evening; you do not need separate lives for glucose, strength, sleep, and peace of mind.

A small pause: Which act of support would help two things at once? A ready lunch might support nourishment and reduce stress. An appointment might bring clarity and stop weeks of guessing.

19. One table, different needs

There is a familiar household sentence: “But what will everyone else eat?”

It can make a personal health goal feel like an additional job. You imagine making two dinners, buying two sets of groceries, and explaining yourself every night. Before long, the work of the plan seems larger than its benefit.

Start with what the people at your table can share.

A protein dish, a vegetable, a sauce, and a carbohydrate component can often be served separately so portions and combinations differ. A taco meal can offer several fillings. A bowl meal can let people choose amounts. A tray of food can become more than one plate without becoming more than one dinner.

The person who wants bread can have bread. The person who needs a different portion can choose it. The person who dislikes mushrooms remains a full member of the household, however mysterious their position may seem to the mushroom enthusiasts.

For couples, a useful conversation begins with help rather than persuasion. “I would like to make lunch more reliable. Could we plan two options together?” Or, “I am trying this dinner pattern. Can we serve the starch separately so we can each build our plate?” You are making a practical request. Your partner does not have to adopt a complete nutritional philosophy before passing the bowl.

For families, avoid making one person's plate the object of constant comment. You can model curiosity, variety, and care without turning the table into a classroom. Children and adults alike deserve a meal without a public assessment of what they chose.

If you cook for one, the shared table may look different, but you still deserve food that feels worth preparing. A meal for one is not a

reduced-value meal. Keep components flexible, use the freezer, and repeat ingredients in ways you enjoy. If loneliness makes eating difficult, consider whether a regular meal with someone, in person or by phone, would be welcome.

Some households need more specialized support. Food allergies, swallowing difficulties, medically prescribed diets, and other needs may limit what can be shared. A dietitian or other qualified professional can help resolve those conflicts. There is no virtue in forcing a general framework into a situation that needs tailored advice.

The practical aim is less duplicated work and more room for ordinary company. Food can support several people without asking them to become the same person.

Notice whether the shared meal is becoming easier to arrange and more pleasant to eat. Less duplicated cooking, fewer explanations, and more ordinary conversation are useful outcomes. They give the food routine a better chance of remaining part of family life, while each person gets the amount and combination that suits them.

At home this week: Choose one meal you already share. Identify which components can stay common and which can be offered separately. Start with a meal people actually like.

20. Restaurants, holidays, and a full life

The invitation arrives, and you want to say yes. Lunch with a friend. A family dinner. A weekend away. Then the food calculations begin.

A useful framework should travel with you lightly enough that you can enjoy where you are going.

At a restaurant, begin with the same questions you use at home. What is the protein? What produce or fiber-containing food is available? What carbohydrate portion fits? What sounds enjoyable? You can ask a practical question about preparation or a side without delivering a speech about your health history.

Whole GI's usual rhythm can guide you, but the occasion may require adaptation. If dinner includes a starch you enjoy, make a thoughtful choice and continue with your normal care. Avoid skipping earlier meals to compensate for a social meal, especially when that conflicts with medication or leaves you overly hungry. Your care plan remains your guide.

Travel adds logistics. Find out where breakfast might come from. Keep a suitable snack available when delays are likely. Plan for medication storage and timing with your care team when necessary. Choose one or two dependable food options near where you are staying. A little preparation can free attention for the trip itself.

At a holiday table, decide what matters to you. There may be a food that carries the whole memory of the day. There may also be several things you eat only because they happen to be there. You can choose with pleasure and awareness. You can have an appropriate portion, enjoy the company, and allow the meal to end.

You do not need to announce that you are being good, bad, careful, or indulgent. "That was lovely" is a complete report.

Alcohol deserves individual guidance, particularly when diabetes medicines are involved; it can increase low-glucose risk.¹⁰ You can enjoy an occasion without it, or follow the advice appropriate to your treatment. There is no requirement to make a health decision for the comfort of everyone else.

You may occasionally encounter someone who wants to comment on your plate. A calm answer can be enough: “I have what I need, thank you.” Then ask about something more interesting. There is usually something more interesting.

When you get home, resume your usual meals. You do not owe the weekend a correction plan. If something was difficult, learn from it: next time a breakfast option, a snack, or a clearer restaurant choice might help. That is adaptation.

A useful sign of progress is being able to enjoy an invitation without spending the whole occasion negotiating with yourself. You can make a considered choice, enjoy the people, and return to your rhythm afterward. The skill is confidence in adapting. The framework is becoming something you can carry with you.

Before your next outing: Choose one practical support in advance. Then give yourself permission to be present for the reason you went.

21. A kitchen that helps you on Tuesday

An organized kitchen can be useful. A kitchen organized for someone else's life can be a very attractive source of frustration.

You need the foods and tools that make your own meals easier.

Begin with the meals you repeat. Look at the ingredients they share. Those are candidates for your dependable supplies: a few protein foods, vegetables in forms you use, fruit you enjoy, suitable carbohydrate options, and flavor ingredients that make everything more appealing. There is no need to buy a complete new pantry in a single trip.

Plan a short distance ahead. Two or three days can be enough when a whole week feels uncertain. Choose meals that use some of the same ingredients so a package does not become an obligation sitting in the refrigerator. Decide where an extra portion will go before you cook it.

Then look at the shopping list against what you already have. Check the freezer. Open the cupboard. The lentils would be delighted to be remembered. Buying an ingredient again because it was hiding behind another one is a remarkably common household subscription service.

Cost matters, and useful nutrition need not depend on premium versions of everything. Compare price per usable serving, consider store brands, and choose convenient forms where they reduce waste or effort. A less expensive ingredient is only a bargain if you can use it. A larger package is only helpful if the extra food has somewhere to go.

Storage and food safety deserve a dependable routine. Refrigerate perishable leftovers promptly, use shallow containers for quicker cooling, and follow safe storage and reheating guidance. USDA recommends refrigerating leftovers within two hours, using refrigerated leftovers within three to four days, and reheating to 165°F.¹⁸ Freeze

portions you will not use in time. Labeling a container is a small act of kindness to the person who will later ask, “What was this?”

The [Pantry Directory](#) offers a place to learn about foods and how they can fit. The [recipe collection](#) can help you turn a few dependable ingredients into different meals. Use these resources when they answer a question, rather than treating them as more material to finish.

The [Meal Planner](#) brings chosen meals and a shopping list together. Adjust servings, consider leftovers, and check what you already have. The value is the work it takes off your mind. A short, usable plan is a good outcome.

The benefit of a dependable kitchen is often felt before the first bite: lunch is available, a backup meal is within reach, and groceries have a purpose. Notice less wasted food, fewer last-minute decisions, and more meals that happen when you need them. Those are the conditions that allow the nutrition principles to become a consistent part of your life.

A small pause: Which ingredient do you repeatedly buy with hopeful intentions and then waste? Choose a more realistic quantity or form next time. Your groceries should fit your habits as they become, not only as you wish them to be.

22. When a difficult week becomes part of the way

You may have a week when the plan barely appears.

Someone gets sick. An appointment becomes a series of appointments. Work expands. Grief arrives. You spend your attention where it is needed, and food becomes whatever is available. Then, when there is a little space again, an old story returns: "I have fallen off."

Before accepting that story, look at what happened. Your circumstances changed. The support available may not have changed with them.

This is where a portable framework proves useful. You can return to one meal without performing a ceremony of starting over. You can use the rescue list. You can ask for help. You can reopen the planner and choose tomorrow's lunch. There is no missed-work penalty attached to the rest of the week.

Begin with the next useful act. Drink as appropriate to your needs. Eat a meal you can manage. Put an easy breakfast within reach. Contact your care team if illness or glucose changes require it. Make the response proportionate to the actual situation.

Then, when you have capacity, ask what would have helped. Perhaps the plan was too complicated even before life became difficult. Perhaps you needed freezer portions, a ready-made option, or someone else to take shopping. Perhaps you need a specific sick-day plan from your clinician. Learning from the week gives you something to carry forward.

Do not make flexibility a prize you receive only after you have followed the plan perfectly. Build it in from the beginning. A routine can have a usual version and an easier version. Both belong to the routine.

This also applies to long transitions. Retirement, a move, bereavement, caregiving, recovery, or a change in household size can alter the

meaning of meals. A former routine may no longer fit. Give yourself room to build a new one without treating the old one as evidence against you.

There may be feelings attached to the kitchen that no grocery list can solve. It is appropriate to seek emotional support as well as practical help. You are a whole person, even in a book about food.

If you are already a member, the resources do not require you to return as an ideal student. Open the part that helps with today's question. A quicksheet, a recipe, or one planned day may be enough to make the next step possible.

One of the most useful changes to notice is a quicker, kinder return. Perhaps a disrupted week once became a month of uncertainty, while now you know which meal to make next. Perhaps you ask for help earlier. That is the framework becoming resilient. Consistency can grow through the ability to adapt and resume.

A small pause: Finish this sentence: "When things become difficult, my easiest useful return is..." Keep the answer small enough that you can believe it.

23. The first thirty days, gently

You do not need a dramatic Monday to begin. You need a first action and a way to repeat it.

For the first week, choose one reliable meal. Breakfast is a useful option if mornings feel uncertain; lunch may be more helpful if the afternoon is difficult. Notice what you currently do, choose a manageable adjustment, and make the ingredients easy to find. The aim is familiarity.

During the second week, connect that meal to another part of the day. If breakfast is becoming dependable, give lunch a proper place. If lunch was your starting point, plan an easier dinner or a useful breakfast. Keep the first change simple while you add the next.

In the third week, practice adaptation deliberately. Use a no-cook option. Share a meal with different portions. Go out and apply the framework without trying to make the experience perfect. Choose one backup meal and test whether it really is easy. An untested rescue meal sometimes turns out to require a food processor and an unexpectedly emotional amount of washing up.

In the fourth week, review what you learned. Which meals were enjoyable enough to repeat? What did you stop buying? Where did you need more support? What became easier? Which medical questions should you bring to your next appointment?

You can slow this down. You can work on one meal for a month. You can begin with access to groceries or a care-team conversation if that is the barrier in front of you. Thirty days is a helpful container for learning, not a deadline for becoming someone else.

Use observations that answer a question. A brief note about lunch, appetite, or sleep may help you notice a pattern. If glucose monitoring has been recommended, follow the schedule and targets you were given.

You do not need to collect every possible metric. Choose information you can use.

An A1C result reflects a longer period, so early progress should not be judged as though a few days can describe the whole picture.² Ask your clinician when to reassess. If readings are concerning or symptoms change, seek guidance rather than waiting for the end of a self-chosen experiment.

You can also notice practical progress: you know what lunch will be, you have an easier dinner available, you are less uncertain about a favorite food. These are meaningful skills even while the medical picture is being followed.

At your review, consider the different time scales. Fullness and satisfaction can change within a meal or a morning. A dependable daily rhythm may become clearer over days and weeks. Strength and clinical measures need their own time and appropriate reassessment. Look for trends rather than demanding that every part of health respond on the same day.

Choose two or three signals that matter most to you: perhaps midmorning cravings, afternoon hunger, and sleep. Make brief notes, then review them alongside any prescribed monitoring. If you feel no clear change yet, that is useful information for adjusting the plan, not a reason to judge your effort. If you do notice improvement, give yourself a moment to recognize it.

Your first page of notes: Write one meal you will practice, one easier backup, and one question for your care team. That is enough to begin a plan.

24. Support you can keep close

By now, you may be able to see your own day in the framework. A breakfast you recognize. A lunch you can anticipate. A dinner that respects the energy you have left. Room for company, a change of plans, and something sweet when you choose it.

You can take what you have learned here into your kitchen today.

If having the pieces together would make it easier, that is what the Well + Easy membership is for. The Low GI Living Meal Planner includes the Whole GI Protocol program, so you can learn the approach and use planning tools alongside it.

The planner helps you choose meals, adjust servings, save favorites, and bring a shopping list together. Cooking Mode offers a way to keep ingredients and instructions in view while you prepare a meal. Preferences and special-diet options can narrow the choices when your circumstances call for it. You remain the person deciding what works in your life.

The program gives you the deeper teaching behind the framework: the workshop, methodology, quicksheets, and resources for applying it, including attention to sleep, stress, inflammation, and hormonal factors. You can use those materials as references. You do not need to complete everything before beginning breakfast.

At the time of this edition, membership is \$15 per month or \$150 per year. Both include the planner and complete Whole GI Protocol program. Plans renew automatically; you can cancel before the next renewal, with access continuing through the paid period. The [membership page](#) has the current details.

I want you to be able to picture the first useful visit. Choose one meal you would enjoy. Put it into a day. Check the ingredients. Decide what you already have and what you need. Let the next action become visible. You can explore further as questions arise.

There are free places to keep learning, too. The recipes, Daily Framework, Pantry Directory, and blog offer different ways into the ideas. The Special Diets collections offer another way to find meals suited to your circumstances. Each link in the web edition opens separately so your place in the book stays here.

You may prefer a screen in the kitchen, a printed page, or a few familiar meals you no longer need to look up. Those are all useful outcomes. The purpose of support is to help you act with more confidence and less unnecessary work.

If you decide to join, come as you are. Bring your usual groceries, your questions, the meals your family loves, and the days when cooking is the last thing you want to do. Those are the materials of a real life.

The membership is there to make these connections easier to use. You can learn the reason behind a practice, choose a meal that applies it, and notice how the routine serves you. Over time, the aim is for the decisions to feel familiar: enough protein, a considered carbohydrate portion, a comfortable evening, and an easier response when plans change. The support helps turn understanding into something you can keep doing.

When you are ready: Explore the Well + Easy membership. Begin with the part that would make tomorrow easier.

25. The life on the other side of the kitchen

An ordinary day can offer a quiet moment of recognition. You make a familiar breakfast without a long debate. You know what lunch will be. Someone invites you out, and the food question feels manageable. Caring for yourself is becoming something you know how to do.

It is a good moment to ask what you want to carry forward.

Perhaps you want to carry less fear about food. Perhaps you want to keep the pleasure of lunch, the comfort of a warm breakfast, or the confidence of having dinner handled. Perhaps you want to carry the knowledge that a difficult week can be met with support instead of a verdict.

You may still have medical questions. You may still want your numbers to change. You may still feel impatient with how much attention a body sometimes requires. Hope does not ask you to pretend those things are gone.

It gives you a reason to take the next useful step while they are here.

You can learn how a favorite food fits. You can make one meal easier. You can bring a clearer question to an appointment. You can make room for enough nourishment, ask for help, or choose a plan that fits the person you are now. These are ordinary actions with room for meaning.

I hope Whole GI gives you a way to connect them. Something clear enough to use when you are busy, flexible enough to accompany change, and kind enough that you can keep returning without rehearsing an apology.

There will still be days when the refrigerator offers ingredients that seem to have met only recently and have no intention of cooperating. There will be meals that do not turn out. You may buy the wrong thing,

forget to defrost something, or enjoy dinner more than you enjoyed planning it. You are allowed to be a person while caring for your health.

And beyond the kitchen, your life is waiting in its own particular shape. A friend to call. A place to visit. Work you care about. A book you keep meaning to finish. A quiet afternoon. People whose company makes the day feel larger.

The food is here to support that life.

Keep the foods you love, with understanding and care. Know one useful thing to do today. Adapt simply when life changes. Let your blood sugar, weight, and strength be part of the picture, while the person you are remains at its center.

You may begin to recognize the changes in small moments: a breakfast that holds you comfortably, fewer insistent cravings, a dinner that leaves you settled, or a night that feels more restful. You may also see meaningful movement in the measures you follow with your care team. Each deserves its proper place. Together, they can help you see the care you are giving your body becoming useful in your life.

You do not have to decide how every future day will go.

Let us make the next meal a little easier, and leave some room for everything that comes after it.

With care,

Jen

NOTES & RESOURCES

Notes and further reading

Whole GI combines clinically studied principles with thirteen years of client application and refinement, as described by its founder. The sources below support the individual practices and findings discussed in the book. Clinical research cited here examines individual practices; the thirteen-year history describes their application and refinement with clients as the Whole GI framework. These are complementary forms of evidence. Observations about client experience, including sleep, are identified as such. Sources were consulted for this edition in October 2026.

1. National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Insulin Resistance & Prediabetes. Overview of insulin resistance and contributing factors. Read the source.
2. NIDDK. The A1C Test & Diabetes. Explains the test, the period it reflects, and factors that may affect interpretation. Read the source.
3. Jenkins DJA, et al. Glycemic index of foods: a physiological basis for carbohydrate exchange. *American Journal of Clinical Nutrition*. 1981;34:362-366. DOI: 10.1093/ajcn/34.3.362. The foundational GI paper. Read the source.
4. Atkinson FS, Brand-Miller JC, Foster-Powell K, Buyken AE, Goletzke J. International tables of glycemic index and glycemic load values 2021: a systematic review. *American Journal of Clinical Nutrition*. 2021;114:1625-1632. DOI: 10.1093/ajcn/nqab233. Values describe tested foods and conditions, not a guaranteed personal response. Read the source.
5. Chiavaroli L, et al. Effect of low glycaemic index or load dietary patterns on glycaemic control and cardiometabolic risk factors in diabetes: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2021;374:n1651. DOI: 10.1136/bmj.n1651. Research on dietary patterns; the average A1C difference is not an individual promise. Read the source.

6. Evert AB, et al. Nutrition Therapy for Adults With Diabetes or Prediabetes: A Consensus Report. *Diabetes Care*. 2019;42:731-754. DOI: 10.2337/dci19-0014. Supports individualized nutrition therapy. [Read the source.](#)
7. Reynolds AN, Akerman AP, Mann J. Dietary fibre and whole grains in diabetes management: systematic review and meta-analyses. *PLOS Medicine*. 2020;17:e1003053. DOI: 10.1371/journal.pmed.1003053. [Read the source.](#)
8. Shukla AP, Iliescu RG, Thomas CE, Aronne LJ. Food Order Has a Significant Impact on Postprandial Glucose and Insulin Levels. *Diabetes Care*. 2015;38:e98-e99. DOI: 10.2337/dc15-0429. A small short-term study; it does not establish universal long-term outcomes. [Read the source.](#)
9. Sutton EF, et al. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes. *Cell Metabolism*. 2018;27:1212-1221.e3. DOI: 10.1016/j.cmet.2018.04.010. A controlled proof-of-concept study with a specific population and schedule. [Read the source.](#)
10. NIDDK. Low Blood Glucose (Hypoglycemia). Medication, meal, activity, and alcohol considerations; treatment requires an appropriate fast-acting carbohydrate plan. [Read the source.](#)
11. Volkert D, et al. ESPEN practical guideline: Clinical nutrition and hydration in geriatrics. *Clinical Nutrition*. 2022;41:958-989. DOI: 10.1016/j.clnu.2022.01.024. Individualized support for adequate nutrition and hydration in older adults. [Read the source.](#)
12. Mozaffarian D, et al. Nutritional Priorities to Support GLP-1 Therapy for Obesity: A Joint Advisory From the American College of Lifestyle Medicine, the American Society for Nutrition, the Obesity Medicine Association, and The Obesity Society. *American Journal of Lifestyle Medicine*. Published online May 30, 2025. DOI: 10.1177/15598276251344827. [Read the source.](#)
13. Deutz NEP, et al. Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. *Clinical Nutrition*. 2014;33:929-936. DOI: 10.1016/j.clnu.2014.04.007. Expert recommendations require individual clinical context. [Read the source.](#)
14. The Menopause Society. Midlife Weight Gain. Patient education on the interacting influences on weight and body composition. [Read the source.](#)

15. Zuraikat FM, et al. Chronic Insufficient Sleep in Women Impairs Insulin Sensitivity Independent of Adiposity Changes: Results of a Randomized Trial. *Diabetes Care*. 2024;47:117-125. DOI: 10.2337/dc23-1156. [Read the source.](#)
16. Reynolds AN, et al. Advice to walk after meals is more effective for lowering postprandial glycaemia in type 2 diabetes mellitus than advice that does not specify timing: a randomised crossover study. *Diabetologia*. 2016;59:2572-2578. DOI: 10.1007/s00125-016-4085-2. [Read the source.](#)
17. Dempsey PC, et al. Benefits for Type 2 Diabetes of Interrupting Prolonged Sitting With Brief Bouts of Light Walking or Simple Resistance Activities. *Diabetes Care*. 2016;39:964-972. DOI: 10.2337/dc15-2336. [Read the source.](#)
18. U.S. Department of Agriculture, Food Safety and Inspection Service. Leftovers and Food Safety. Practical storage and reheating guidance. [Read the source.](#)
19. National Center for Complementary and Integrative Health (NCCIH). Chamomile. Safety information includes potential interactions with medicines. [Read the source.](#)
20. Dalgaard LB, et al. A dairy-based, protein-rich breakfast enhances satiety and cognitive concentration before lunch in overweight to obese young females: A randomized controlled crossover study. *Journal of Dairy Science*. 2024;107:2653-2667. DOI: 10.3168/jds.2023-24152. Greater short-term fullness did not translate into lower total daily calories in this study. [Read the source.](#)
21. Leidy HJ, et al. Beneficial effects of a higher-protein breakfast on the appetitive, hormonal, and neural signals controlling energy intake regulation in overweight/obese, breakfast-skipping, late-adolescent girls. *American Journal of Clinical Nutrition*. 2013. DOI: 10.3945/ajcn.112.053116. This population-specific study informs appetite guidance, rather than predicting an identical response at every age. [Read the source.](#)
22. Gu C, et al. Metabolic Effects of Late Dinner in Healthy Volunteers - A Randomized Crossover Clinical Trial. *Journal of Clinical Endocrinology & Metabolism*. 2020;105:2789-2802. DOI: 10.1210/clinem/dgaa354. Compared dinner at 10 p.m. and 6 p.m.; the finding supports attention to timing and is not a trial of Whole GI dinner composition. [Read the source.](#)
23. Jakubowicz D, et al. High-energy breakfast with low-energy dinner decreases overall daily hyperglycaemia in type 2 diabetic patients: a randomised clinical trial. *Diabetologia*. 2015;58:912-919. DOI: 10.1007/s00125-015-3524-9. Examines redistribution of food earlier in the day; it did not test the exact Whole GI schedule. [Read the source.](#)

24. Vlahoyiannis A, et al. A Systematic Review, Meta-Analysis and Meta-Regression on the Effects of Carbohydrates on Sleep. *Nutrients*. 2021;13:1283. DOI: 10.3390/nu13041283. Different carbohydrate intakes affected different aspects of sleep; this does not establish that a non-starchy dinner universally improves sleep. Sleep benefits described from client experience are observations about the broader routine. [Read the source.](#)
25. NIDDK. Eating, Diet, & Nutrition for Constipation. Fiber, fluids, and bowel regularity. [Read the source.](#)
26. NIDDK. Eating, Diet, & Nutrition for GER & GERD. Includes timing advice for nighttime reflux symptoms. [Read the source.](#)
27. Verreijen AM, et al. A higher protein intake at breakfast and lunch is associated with a higher total daily protein intake in older adults: a post-hoc cross-sectional analysis of four randomised controlled trials. *Journal of Human Nutrition and Dietetics*. 2021;34:384-394. DOI: 10.1111/jhn.12838. An association in older adults, rather than a direct test of muscle gain. [Read the source.](#)
28. Morton RW, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *British Journal of Sports Medicine*. 2018;52:376-384. DOI: 10.1136/bjsports-2017-097608. [Read the source.](#)
29. Zhang J, et al. Effects of Water Restriction and Supplementation on Cognitive Performances and Mood among Young Adults in Baoding, China: A Randomized Controlled Trial (RCT). 2021. Examines rehydration after water restriction; morning water is the framework's practical cue for hydration. [Read the source.](#)
30. Drake C, et al. Caffeine effects on sleep taken 0, 3, or 6 hours before going to bed. *Journal of Clinical Sleep Medicine*. 2013. Examines caffeine and sleep; supports choosing caffeine-free evening drinks. [Read the source.](#)
31. Estruch R, et al. Primary Prevention of Cardiovascular Disease with a Mediterranean Diet Supplemented with Extra-Virgin Olive Oil or Nuts. *New England Journal of Medicine*. 2018;378:e34. The corrected, republished PREDIMED analysis. [Read the source.](#)
32. Sonia S, et al. Effect of cooling of cooked white rice on resistant starch content and glycemic response. *Asia Pacific Journal of Clinical Nutrition*. 2015;24:620-625. DOI: 10.6133/apjcn.2015.24.4.13. Food variety, portion, and safe handling remain part of applying the finding. [Read the source.](#)

33. Surwit RS, et al. Stress management improves long-term glycemic control in type 2 diabetes. *Diabetes Care*. 2002;25:30-34. DOI: 10.2337/diacare.25.1.30. A structured program followed over a year, rather than a test of a single brief relaxation practice. Read the source.

34. Esposito K, et al. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA*. 2004;292:1440-1446. DOI: 10.1001/jama.292.12.1440. Read the source.

35. Ortinau LC, et al. Effects of high-protein vs. high-fat snacks on appetite control, satiety, and eating initiation in healthy women. *Nutrition Journal*. 2014. Examines specific afternoon snacks in young adult women; composition, portion, and individual needs guide practical use. Read the source.

Keep exploring

The book is a place to return to. These resources can help you put a chapter into practice.

- **Membership and current pricing:** the Low GI Living Meal Planner and included Whole GI Protocol program.
- **The Daily Framework:** a morning-to-night overview.
- **Recipes:** familiar meals to explore.
- **Pantry Directory:** learn about ingredients and how they can fit.
- **Special Diets:** browse options suited to different preferences and circumstances.
- **Whole GI member resources:** program lessons and downloads; membership and sign-in are required.
- **Whole GI Protocol overview:** learn about the program behind the planner.
- **Well + Easy blog:** articles and further learning.

© 2026 Well + Easy. The Whole GI Way. Revised edition, 2026.
Educational nutrition and lifestyle guidance; individual medical advice takes priority.